

Monthly report tables and figures (April participants)

Figure 1. Participants recruited in April by municipality of residence (N=887)




Table 1. Distribution of participants and COVID-19 cases reported in April by participants for last 30 days, by age and gender

	Participants	Confirmed cases (RT-PCR)	Confirmed and probable cases*
	Number (column%)	Number (row%)	
Number	887	12 (1.4)	38 (4.3)
Sex			
Male	275 (31.0)	3 (1.1)	13 (4.7)
Female	598 (67.4)	9 (1.5)	25 (4.2)
Missing/Other	14 (1.6)	0 (0)	0 (0)
Age			
21-25	40 (4.7)	1 (2.5)	2 (5.0)
26-30	43 (5.1)	0 (0)	3 (7.0)
31-35	45 (5.3)	0 (0)	2 (4.4)
36-40	65 (7.7)	3 (4.6)	6 (9.2)
41-45	91 (10.8)	0 (0)	1 (1.1)
46-50	99 (11.7)	2 (2.0)	4 (4.0)
51-55	112 (13.2)	2 (1.8)	8 (7.1)
56-60	114 (13.5)	1 (0.9)	6 (5.3)
61-65	91 (10.8)	0 (0)	0 (0)
≥66	146 (17.3)	2 (1.4)	5 (3.4)

Cases percentages computed as a percent of participants in each age and sex category

*Confirmed and probable cases (including other tests and/or model based on key symptoms) [4]

The numbers may not add to 887 due to missing data.

Table 2. Status of COVID-19 cases reported in April for last 30 days

	Confirmed cases (RT-PCR) N=4	Confirmed and probable cases* N=16
	Number (%)	
Cases with symptoms \geq 21 days	0 (0)	3 (18.8)
Cases admitted to the hospital	1 (25.0)	1 (6.3)
Cases admitted to the Intensive Care Unit (ICU)	0 (0)	0 (0)
Cases who received oxygen treatment	0 (0)	0 (0)

Not all cases answer these questions, therefore percentages may not be representative of the total cases.

*Confirmed and probable cases (including other tests and model based on key symptoms)^[4]

Table 3. Percent of people who reported specific symptoms in the last 30 days

	Overall	Within confirmed COVID-19 cases (RT-PCR)	Within all COVID-19 cases*
	Number (%) with symptoms		
Number of people	887	12	38
Symptoms			
Fever	45 (5.1)	3 (25.0)	14 (36.8)
Nausea/vomiting	52 (5.9)	2 (16.7)	13 (34.2)
Diarrhea	76 (8.6)	3 (25.0)	14 (36.8)
Persistent cough	34 (3.8)	2 (16.7)	11 (29.0)
Difficulty breathing	49 (5.5)	1 (8.3)	12 (31.6)
Unusual fatigue	70 (7.9)	2 (16.7)	17 (47.2)
Loss of taste	29 (3.3)	3 (25.0)	22 (57.9)
Loss of smell	29 (3.3)	3 (25.0)	22 (57.9)
Chest discomfort, tension, pain and/or pressure	65 (7.3)	4 (33.3)	17 (44.7)
Skipping meals	128 (14.5)	3 (25.0)	18 (47.4)
Delirium	25 (2.8)	0 (0)	6 (15.8)
Muscle and/or joint pain	210 (23.7)	4 (33.3)	24 (63.2)

*Confirmed and probable cases (including other tests and model based on key symptoms)^[4]

The number of cases may not add to the confirmed or total cases due to missing data.

Table 4. COVID-19 cases by key comorbidities

	Confirmed COVID-19 cases (RT-PCR)	Total COVID-19 cases*
	Number (%)	
With Diabetes (N=150)	2 (1.3)	5 (3.3)
Without Diabetes (N=718)	10 (1.4)	31 (4.3)
With Pulmonary conditions (including asthma) (N=118)	3 (2.5)	7 (5.9)
Without pulmonary conditions (N=748)	9 (1.2)	29 (3.4)
Obese** (N=399)	6 (1.5)	19 (4.8)
Not Obese (N=444)	6 (1.4)	19 (4.3)

*Confirmed and probable cases (including other tests and model based on key symptoms)^[4]

**Participants were classified as obese if they have a body mass index (BMI) ≥ 30

The number of cases may not add to the confirmed or total cases due to missing data

Figure 2. Monthly COVID-19 positive cases reported in April for the last 30 days (Confirmed cases by RT-PCR)




Figure 3. Monthly COVID-19 positive cases reported in April for the last 30 days (All confirmed and probable cases including other tests and model based on key symptoms)




Table 5. Influenza positive cases reported in April for the last 30 days among those who reported being tested in the last 30 days (N=103)

	Participants	Tested for Influenza	Influenza cases
	Number (row %)		
Number	887	103 (11.6)	3 (0.3)
Sex			
Male	275 (31.0)	36 (13.1)	1 (0.4)
Female	598 (67.4)	65 (10.9)	2 (0.3)
Missing/Other	14 (1.6)	2 (14.3)	0 (0)
Age			
21-25	40 (4.7)	6 (15.0)	0 (0)
26-30	43 (5.1)	1 (2.3)	0 (0)
31-35	45 (5.3)	5 (11.1)	1 (2.2)
36-40	65 (7.7)	8 (12.3)	0 (0)
41-45	91 (10.8)	10 (11.0)	0 (0)
46-50	99 (11.7)	10 (10.1)	1 (1.0)
51-55	112 (13.2)	14 (12.5)	0 (0)
56-60	114 (13.5)	15 (13.2)	0 (0)
61-65	91 (10.8)	16 (17.6)	0 (0)
≥66	146 (17.3)	14 (9.6)	0 (0)

Cases percentages computed as a percent of participants in each age and sex category

Figure 4. Monthly Influenza positive cases reported in April for the last 30 days




Table 6. COVID-19 vaccination distribution by age range and comorbidities

	Participants	Only one dose	Two doses
	Number (%)		
Number	887	234 (26.9)	439 (50.5)
Age			
21-25	40 (4.7)	10 (25.6)	12 (30.8)
26-30	43 (5.1)	7 (16.7)	16 (38.1)
31-35	45 (5.3)	12 (26.7)	14 (31.1)
36-40	65 (7.7)	14 (21.5)	24 (36.9)
41-45	91 (10.8)	41 (46.1)	29 (32.6)
46-50	99 (11.7)	35 (36.5)	40 (41.7)
51-55	112 (13.2)	35 (32.1)	51 (46.8)
56-60	114 (13.5)	37 (32.5)	58 (50.9)
61-65	91 (10.8)	25 (27.8)	56 (62.2)
≥66	146 (17.3)	10 (7.1)	122 (87.1)
Comorbidities			
With Diabetes	150 (16.9)	41 (27.5)	88 (59.1)
With Pulmonary conditions (including asthma)	118 (13.3)	39 (33.3)	55 (47.0)
Obese	399 (44.9)	116 (29.6)	186 (47.5)

The number of people who responded COVID-19 vaccination question was 869

Table 7. Requirement to work close to other people during the pandemic by COVID-19 vaccination status

	Not vaccinated	Vaccinated (one dose)	Vaccinated (two doses)
	Number (%)		
Required to work close to others			
Yes	17 (73.9)	17 (56.7)	38 (76.0)
No	6 (26.1)	13 (43.3)	12 (24.0)

The number of people who responded to this module and vaccine question was 103

Table 8. Willingness to get vaccinated against COVID-19

	No	Yes	Not sure
	Number (%)		
Number	3 (15.0)	11 (55.0)	6 (30.0)
Age			
21-25	0 (0)	3 (75.0)	1 (25.0)
26-30	0 (0)	1 (100)	0 (0)
31-35	0 (0)	1 (50.0)	1 (50.0)
36-40	1 (25.0)	2 (50.0)	1 (25.0)
41-45	0 (0)	0 (0)	0 (0)
46-50	0 (0)	2 (100)	0 (0)
51-55	0 (0)	2 (50.0)	2 (50.0)
56-60	0 (0)	0 (0)	0 (0)
61-65	0 (0)	0 (0)	0 (0)
≥66	1 (50.0)	0 (0)	1 (50.0)
Comorbidities			
With Diabetes	0 (0)	1 (100)	0 (0)
With Pulmonary conditions (including asthma)	0 (0)	2 (66.7)	1 (33.3)
Obese	1 (12.5)	3 (37.5)	4 (50.0)
Nature of work			
Not currently working	1 (9.1)	8 (72.7)	2 (18.2)
Healthcare sector	0 (0)	0 (0)	0 (0)
An essential worker, not in health sector	1 (16.7)	2 (33.3)	3 (50.0)
Not from the health sector nor an essential worker	1 (33.3)	1 (33.3)	1 (33.3)

The number of people who responded to this module and was not yet vaccinated for COVID-19 was 20

Figure 5. Comparison of COVID-19 and Influenza cases in the last 30 days from November 2020 to April 2021




Figure 6. Comparison of the weeks of April 2021 for COVID-19 cases in the last 30 days

